




# Continuous gravitational wave atlas.


Vladimir Dergachev  
Max Planck/AEI Hannover



Anaheim, 2025-Mar-17

# Somewhere far away there is a neutron star...

We want to find it !



Bump  
(not to  
scale)

Circularly polarized  
gravitational waves


- Hard, computationally intensive problem
- Small parameter – equatorial deformation of neutron star  $\epsilon$
- Sensitivity scales as  $(\text{coherence length})^{-0.25}(\text{frequency})^2$  and is proportional to  $\epsilon$
- Computing time scales as  $(\text{coherence length})^4(\text{frequency})^3$  or faster

Linearly  
polarized  
gravitational  
waves


# Continuous gravitational waves

- Need a rotating star with **non-zero equatorial second moment**
- Gravitational radiation is expected to be emitted at **twice the rotation frequency**
- Continuous wave signals have very **narrow bandwidth**
- The only signal that can be measured **again**, months and years after detection

Rotating neutron star




Linearly  
polarized  
gravitational  
waves



Circularly polarized  
gravitational waves


# Atlas as a transform of detector data

- The detectors produce a time series of strain values.
- It is relatively easy to do a search for short transient signals in time series.
- But it is very difficult to search for continuous signals – you have go through entire data for every sky location, every frequency and every frequency derivative.
- It would be easier to work with data already in frequency domain.
- This is an atlas of continuous gravitational waves - data over sky and frequency !



# Continuous gravitational wave atlas


- Instant search on your notebook !
- Robust results for the entire sky and all covered frequencies, no exclusions.
- First atlas released 2022-02-22, covering 500-1000 Hz



Phys. Rev. X 13, 021020 (2023)

# Several atlas releases available


- First atlas released 2022-02-22, covering 500-1000 Hz,  
[Phys. Rev. X 13, 021020 \(2023\)](#)
- Expanded early release atlas released 2023-11-16, covering 20-1500 Hz and larger spindown range, [Phys. Rev. D 109, 022007 \(2024\)](#)
- **Just released:** Full 20-1700 Hz atlas [arXiv:2503.11512](#)
- Binary spotlight search atlas [arXiv:2503.11503](#)
- There is also Gaia DR3 data in the same format as the atlas for easy analysis
- Each data set has examples of common searches





# Continuous gravitational wave atlas


- An atlas is a collection of sky maps, such as shown on the right
- Separate sky maps for each 45mHz frequency
- Each pixel has data for many different metrics



# Metrics

Each sky location and frequency band has the following metrics:


- maximum SNR
- frequency and polarization where SNR was achieved
- upper limit on arbitrary polarized signals (“worst case”)
- upper limits for circularly polarized signals
- data to compute polarization specific upper limits





# Atlas is constructed by analyzing signal power


- The  $h(t)$  data is demodulated, passed through narrow band filter, squared and summed.
- This makes it easy to relate the output statistics with the strength of astrophysical signals
- Plot on the right shows upper limits, derived using power measurements.
- Longer coherence length (12 hours+) makes data more sensitive than LVK
- Use atlas now, no need to wait for O4 results



# The power is analyzed using universal statistics


- Upper limits are produced using universal statistic method
- Universal means that it does not assume any particular distribution of noise in input data.
- The penalty is a slight overestimate of upper limits, compared to what could be achieved if distribution was known (plot on the right)
- Atlas reports 95% confidence level upper limits. They are valid in all frequency bands, over all the sky – no exclusions

Phys. Rev. D 87, 062001 (2013)




# Usage examples: interactive browser

- R example `view_summary.R`
- Interactively displays maximums of SNR or upper limits across a frequency band and over sky
- Plot on the right was made with `plot_gw(835, 840, "snr")`
- Also see `view_help()`



# Usage examples: investigating single templates

- R example  
spatial\_index\_example1.R
- Uses spatial index to quickly find data for specific sky location and frequency



```
1 library("RMVL")
2 #
3 # In this example we use an input file "points_of_interest.csv" which contains
4 # coordinates for several points of interest (f0, ra, dec)
5 # We use previously constructed spatial index to find points in the O3a atlas that are
6 # closest to points of interest
7 # Results are printed in the end of the file
8 #
9 # O3a_atlas.mvl contains results of O3a analysis
10 Matlas<-mvl_open("O3a_2_atlas.mvl")
11
12 # O3a_spatial_index.mvl is created by script create_spatial_index.R from O3a_atlas.mvl
13 Mspatial<-mvl_open("O3a_2_spatial_index.mvl")
14
15 skymaps_band_start<-Mspatial$skymaps_band_start
16 x0<-Mspatial$x0
17 x1<-Mspatial$x1
18 x2<-Mspatial$x2
19
20 # User provided files. Example included.
21 # For very large files, it might be better to store data in MVL format
22 points_of_interest<-read.table("points_of_interest.csv", header=TRUE)
23
24 f0poi<-points_of_interest[, "frequency"]
25 x0poi<-cos(points_of_interest[, "ra"])*cos(points_of_interest[, "dec"])*f0poi/7.5
26 x1poi<-sin(points_of_interest[, "ra"])*cos(points_of_interest[, "dec"])*f0poi/7.5
27 x2poi<-sin(points_of_interest[, "dec"])*f0poi/7.5
```

# Usage examples: investigating specific sky location


- R example  
spatial\_index\_example3.R
- Uses spatial index to quickly find all data for a specific sky location

```
spatial_index_example3.R — KWrite
File Edit View Bookmarks Tools Settings Help
New Open... Save Save As... Close Undo Redo

1 library("RMVL")
2 #
3 # In this example we retrieve upper limits and signal to noise ratios for a particular
4 sky location, using previously constructed spatial index.
5 #
6 # 03a_atlas.mvl contains results of 03a analysis
7 Matlas<-mvl_open("03a_2_atlas.mvl")
8
9 # 03a_spatial_index.mvl is created by the script create_spatial_index.R from
10 03a_atlas.mvl
11 Mspatial<-mvl_open("03a_2_spatial_index.mvl")
12
13 skymaps_band_start<-Mspatial$skymaps_band_start
14 x0<-Mspatial$x0
15 x1<-Mspatial$x1
16 x2<-Mspatial$x2
17
18 # Coordinates in radians, J2000
19 #
20 RA<-0.0
21 DEC<-0.0
22
23 # Search all frequencies
24 f0poi<-sort(unique(Matlas$parameters[, "band_start"][]))
25
26 # Rescale for spatial index
27 x0poi<-cos(RA)*cos(DEC)*f0poi/7.5
28 x1poi<-sin(RA)*cos(DEC)*f0poi/7.5
29 x2poi<-sin(DEC)*f0poi/7.5
```

# Example: directed searches

- G189.1+3.0 – data on the right is an extract from Falcon atlas
- Latest LVK results shown for comparison





# Summary

- Atlas provides all-sky, spectrally resolved data for continuous gravitational wave sources – a starting point for new searches
- New MVL file format for large scale data analysis
- Ready to use examples of searches using Falcon atlas releases and Gaia DR3 data
- Get the data:

<https://www.atlas.aei.uni-hannover.de/work/volodya/releases.html>

END OF TALK

# Polarization specific upper limits


- Atlas includes *functional* upper limits, with a separate 95% confidence level value for each  $i$  and  $\psi$
- The coefficients  $c_1$ - $c_{14}$  are chosen large enough that upper limits are always valid
- It is possible to choose them so that the overestimate is small ( $\sim 5\%$ ) for noise dominated data

$$\widehat{\text{UL}}^2 = (c_1 + f_{pp}c_2 + f_{pc}c_3 + f_{cc}c_4 + f_{impc}c_5 + f_{pp}^2c_6 + f_{cc}^2c_7 + f_{pc}^2c_8 + f_{impc}f_{pp}c_9 + f_{impc}f_{pc}c_{10} + f_{impc}f_{cc}c_{11} + f_{pp}f_{pc}c_{12} + f_{cc}f_{pc}c_{13} + f_{pp}f_{cc}c_{14}) / (f_{pp} + f_{cc})$$

$$\begin{aligned} a_+ &= \frac{(1 + \cos^2 \iota)^2}{4} \\ a_\times &= \cos^2 \iota \\ f_{pp} &= 2|\tilde{w}_1|^2 = \frac{1}{4} (a_+ + a_\times + (a_+ - a_\times) \cos 4\psi) \\ f_{pc} &= 4\text{Re} \tilde{w}_1 \tilde{w}_2^* = \frac{1}{2} ((a_+ - a_\times) \sin 4\psi) \\ f_{cc} &= 2|\tilde{w}_2|^2 = \frac{1}{4} (a_+ + a_\times - (a_+ - a_\times) \cos 4\psi) \\ f_{impc} &= 2\text{Im} \tilde{w}_1 \tilde{w}_2^* = \frac{1}{4} (1 + \cos^2 \iota) \cos \iota \end{aligned}$$

# Atlas data uses MVL file format

- Designed for efficient access by memory mapping
- Useful for interactive and scripted analysis of large data
- Example plot on the right was constructed using Gaia data in MVL format



# MySQL/MariaDB/sqlite vs MVL files

Mysql/MariaDB/sqlite/Postgresql	MVL
Collection of tables, each table consists of fixed length rows	Can store tables, but also lists, trees and other complex data structures
Lookup based indices, usually $\log(N)$ scaling	Hash based indices – $O(1)$ scaling with length
	Spatial indices - find objects near query
Needs setup, dedicated server	Just files – use as is
Server needs to be large enough to support cluster usage	Files are memory mapped and just need a fast enough file system.
	Loaded data is shared between processes
Supports bulk data storage as well as frequently changed data, such as created by transactions	Focused on large data storage, optimized for solid state drives

# Example: hardware injections

TABLE I. This table shows parameters of the hardware-injected continuous wave signals and atlas data for their locations and frequencies. The upper limits for the injections are polarization specific and were computed using  $\iota$  and  $\psi$  of each injection. We show all the hardware injections within 20–1500 Hz range, including those outside of our search space, as indicated by the “In” column. We use the reference time (GPS epoch)  $t_0 = 1246070000$  (2019 Jul 2 02:33:02 UTC).


Label	$f$ Hz	$\dot{f}$ Hz/s	Binary	SNR	UL/ $h_0$ %	$\Delta f$ mHz	In
ip0	265.57505	$-4.15 \times 10^{-12}$	No	28.5	122.5	−0.1	Yes
ip1	848.93498	$-3 \times 10^{-10}$	No	393.0	119.9	−0.1	Yes
ip2	575.16351	$-1.37 \times 10^{-13}$	No	39.3	138.5	0.0	Yes
ip3	108.85716	$-1.46 \times 10^{-17}$	No	23.7	141.6	0.1	Yes
ip4	1390.60583	$-2.54 \times 10^{-8}$	No	7.6	21.3	−7.7	No
ip5	52.80832	$-4.03 \times 10^{-18}$	No	155.9	130.2	0.0	Yes
ip6	145.39178	$-6.73 \times 10^{-9}$	No	8.4	25.0	−11.2	No
ip7	1220.42586	$-1.12 \times 10^{-9}$	No	7.3	68.1	3.6	No
ip8	190.03185	$-8.65 \times 10^{-9}$	No	8.9	83.8	−2.9	No
ip9	763.84732	$-1.45 \times 10^{-17}$	No	39.1	135.1	0.1	Yes
ip10	26.33210	$-8.5 \times 10^{-11}$	No	63.9	124.9	0.0	Yes
ip11	31.42470	$-5.07 \times 10^{-13}$	No	93.2	400.9	−12.1	Yes
ip12	37.75581	$-6.25 \times 10^{-9}$	No	14.0	156.5	4.0	No
ip16	234.56700	0	Yes	8.3	29.6	42.7	No
ip17	890.12300	0	Yes	8.1	103.6	23.6	No



# Continuous gravitational wave atlas

- New early release (2023 Nov) while we are still analyzing outliers
- 20-1500 Hz
- $|\dot{f}| < 5e-10$  Hz/s
- Data from two stages, 12 and 24 hour coherence length

Phys. Rev. D 109, 022007 (2024)



# How to make an all-sky search


- Partition the sky into small patches. Partition frequency into small bands. Partition any other parameters like frequency directives.
- The neutron star you are looking for can be in any of these patches of parameter space.
- Write code that reliably detects neutron star in a patch and rejects detector noise.
- Rejection of detector noise typically implies that you reject neutron stars in patches you are not searching.
- Now you need to iterate your search over all the patches.
- If you have a patch where you have not detected a strong signal you could analyze it with a more sensitive code, or place an upper limit.
- There are too many patches – in the past we would just take the maximum over all parameters except frequency.

# Falcon – Fast Loosely Coherent Search

- Designed for wide band all-sky searches
- Optimized for analysis with coherent lengths from few hours to several days.
- Worst case upper limits are computed as maximum over sky and frequency derivative. They are valid for any subset
- Detection pipeline produces high quality outliers



Circularly polarized  
gravitational waves




Linearly  
polarized  
gravitational  
waves


Phys. Rev. Lett. 123, 101101 (2019)  
Phys. Rev. D 101, 022001 (2020)  
Phys. Rev. Lett. 125, 171101 (2020)  
Phys. Rev. D 103, 063019 (2021)  
Phys. Rev. X 13, 021020 (2023)

# What is a loosely coherent search ?


Conventional matched filter looks for one waveform at a time.  
Sensitive, but very large parameter space



Semi-coherent searches partition data and integrate results of analysis in each chunk. Sensitivity lost due to **unphysical waveforms**.



Loosely coherent search analyses sets of trajectories at a time. The set of allowed waveforms is controlled for best sensitivity and computational efficiency



# Low-ellipticity pulsars

- It is known that neutron star crust can support ellipticities of  $\approx 10^{-6}$
- But we do not know what physical process will produce them naturally
- No detections in previous searches
  - This might be due to lack of sensitivity, with signals just below noise floor
  - Or because natural sources do not perfectly follow assumed model

*There are generic arguments that many known pulsars have ellipticities of  $10^{-8}$  and that there is a minimum ellipticity of  $10^{-9}$*

[ApJ 863 2](#) G. Woan, M. D. Pitkin, B. Haskell, D. I. Jones, P. D. Lasky

# Low-ellipticity pulsars

- Plot on the right shows distance to pulsars with ellipticity of  $10^{-8}$
- We are sensitive to sources up to 150 pc away
- Frequency derivatives up to  $\pm 5 \cdot 10^{-11}$
- +50% sensitivity compared to O2

